

## Complete DAGs

[DAG is just another TLA I don't approve of; it stands for "Directed Acyclic Graph".]

In a recent manuscript - in the literal sense of the word - "Some properties of the relative converse" (March 1995), Tony Hoare plays with tetrahedra of which the edges may be directed. (As by-product of his considerations he finds, for instance, that "any non-cyclic ascription of directions to any five of the edges can be non-cyclically extended to the sixth edge.") Here is a little bit of related theory.

\*            \*            \*

Let  $K$  be the complete graph with  $n$  vertices, in which each edge has a direction. Then the following 3 statements are equivalent

- (i)  $K$  contains no cyclic paths
- (ii)  $K$  contains no cyclic paths of length 3
- (iii) the outdegrees of the vertices of  $K$  are the values 0 through  $n-1$

The equivalences hold for  $n=1$ , since

for  $n=1$  (i), (ii), and (iii), are all true. For general  $n$ , the proof proceeds by mathematical induction. The induction step itself is done by cyclic implication: we now consider the complete  $(n+1)$ -graph  $K'$  with directed edges, and observe for it

(i)  $\Rightarrow$  (ii)

This follows -independently of the induction hypothesis- from the fact that "of length 3" is a restriction.

(ii)  $\Rightarrow$  (iii)

Let  $A$  be of  $K'$  a vertex of maximum outdegree; let  $K$  be the  $n$ -graph that remains after removal of  $A$  and its  $n$  connecting edges. Because of (ii), also  $K$  contains no cycles of length 3 and, ex hypothese, the outdegrees of its vertices are the values 0 through  $n-1$ ; consequently we are done when we show that  $A$  has outdegree  $n$  (note that then the vertices of  $K$  have the same outdegree in  $K$  as in  $K'$ ).

Let  $B$  be the vertex with outdegree  $n-1$  in  $K$ ; since, by construction,  $A$  is not a vertex of  $K$ ,  $A$  and  $B$  are different vertices. We now first deal with

with the edge  $AB$ , and then with the remaining edges connecting  $A$  to  $K$ .

The direction of edge  $AB$  is  $A \rightarrow B$  for the assumption  $B \rightarrow A$  leads to a contradiction: then  $B$  would have in  $K'$  the (maximum) outdegree  $n$ , and so would  $A$  (by virtue of how it has been chosen: outdegree  $A \geq$  outdegree  $B$ ), but in the directed complete  $(n+1)$ -graph, at most 1 vertex has the maximum outdegree  $n$ .

Let  $C$  be a third vertex; then, because  $C$  is a vertex of  $K$ , in which  $B$  has the maximal outdegree  $n-1$ , the direction of edge  $BC$  is  $B \rightarrow C$ . From  $A \rightarrow B$ ,  $B \rightarrow C$  and the absence of cycles of length 3 in  $K'$ , we conclude that the direction of  $AC$  is  $A \rightarrow C$ . (Note that, so far, we had only used the absence of cycles in  $K$ .) Hence,  $A$  has outgoing edges only, quod erat demonstrandum.


(iii)  $\Rightarrow$  (i)

Assume (iii) for  $K'$ ; let  $A$  be the vertex of maximum outdegree  $n$ ; let  $K$  be the  $n$ -graph that remains after removal of  $A$  and its  $n$  connecting

edges. Possible cyclic paths in  $K'$  are then of two kinds, either they include  $A$ , or they lie in  $K$ .

Because  $A$  has outgoing edges only, no cyclic path leads through it; because of assumption (iii) for  $K'$  and of the fact that  $A$  has outgoing edges only, we conclude (iii) for  $K$  and, ex hypothesi, that no cyclic paths lie in  $K$ . So,  $K'$  contains no cyclic paths, quod erat demonstrandum.

\* \* \*

The absence of cyclic paths in the complete graph tells us - with apologies for the picture! - that each triangle is of the form , i.e.  $\rightarrow$  is a transitive relation. Hiding this fact so far, could be considered a conscious obfuscation on my part.

Austin, 29 March 1995

prof. dr. Edsger W. Dijkstra  
 Department of Computer Sciences  
 The University of Texas at Austin  
 Austin, TX 78712-1188  
 USA