

1. Present a combinatorial argument that for all positive integers  $x$  and  $y$

$$\sum_{k=0}^n \binom{n}{k} x^k y^{n-k} = (x+y)^n.$$

(Hint: Consider sequences drawn from the union of distinct sets  $A$  and  $B$  of cardinalities  $x$  and  $y$ , respectively.)

2. Present a combinatorial argument that for all positive integers  $1 \leq k \leq m \leq r$ :

$$\binom{r}{m} \binom{m}{k} = \binom{r}{k} \binom{r-k}{m-k}.$$

3. a. Present a combinatorial argument that for all positive integers  $n$ ,  $a$ , and  $b (>a)$ :

$$\sum_{k=0}^n \binom{n}{k} a^k (b-a)^{n-k} = b^n.$$

- b. Present a combinatorial argument that for all positive integers  $n$ :

$$\binom{2n}{2} = 2 \binom{n}{2} + n^2$$

4. Using a combinatorial argument, prove that for  $n \geq 1$ :

$$\sum_{k=0}^n \binom{n}{k}^2 = \binom{2n}{n}$$

5. a. Present a combinatorial argument that for all positive values of  $n$ :

$$3^n = \sum_{i=0}^n \sum_{j=0}^{n-i} \binom{n}{i \quad j}$$

- b. Present a combinatorial argument that for all  $m$  and  $n$  satisfying  $2 \leq m$ ,  $2 \leq n$ , and  $m \leq n+1$ :

$$\binom{n+2}{m} = \binom{n+1}{m} + \binom{n}{m-1} + \binom{n}{m-2}$$

(Hint: Consider  $A = B \cup \{c\} \cup \{d\}$ , where  $c \neq d$ ,  $c \notin B, d \notin B$ , and  $\#B = n$ .)

6. a. Present a combinatorial argument that for all  $n$  and  $k$  satisfying  $1 \leq n$  and  $k \leq n$ :

$$n! = \binom{n}{k} \cdot k! \cdot (n-k)!$$

- b. Present a combinatorial argument that for all positive values of  $n$ :

$$2^n = 1 + \sum_{k=0}^{n-1} 2^k$$

(Hint: Consider Let  $k$  be the position of the first 1 in a bit string.)

7. a. Present a combinatorial argument that for all  $n \geq 1$ :

$$\sum_{k=0}^n \binom{n}{k} 2^k = 3^n$$

b. Present a combinatorial argument that for all nonnegative integers  $p, s$ , and  $n$  satisfying  $p + s \leq n$

$$\binom{n}{p} \binom{n-p}{s} = \binom{n}{p+s} \binom{p+s}{p}$$

(Hint: Consider choosing two subsets.)

8. a. Present a combinatorial argument that for all  $n \geq 1$ :

$$\sum_{k=1}^n \binom{n}{k} = 2^n - 1$$

(Note: The summation begins with  $k = 1$ .)

b. Present a combinatorial argument that for all integers  $k$  and  $n$  satisfying  $3 \leq k \leq n$

$$\binom{n}{k} = \binom{n-3}{k} + 3 \binom{n-3}{k-1} + 3 \binom{n-3}{k-2} + \binom{n-3}{k-3}$$

(Hint: Consider three special elements.)

9. Present a combinatorial argument that for all positive integers  $m, n$ , and  $r$ , satisfying  $r \leq \min\{m, n\}$ :

$$\binom{m+n}{r} = \sum_{k=0}^n \binom{m}{k} \binom{n}{r-k}$$

(Hint: Consider selecting from two sets.)

b. Present a combinatorial argument that for all positive integers  $n$ :

$$3^n = \sum_{i=0}^n \left( \sum_{j=0}^{n-i} \binom{n}{i} \binom{n-i}{j} \right)$$

(Note: Be very specific about the roles of  $i$  and  $j$ .)