

**M 340L - CS**  
**Homework Set 8**

1. Let  $u^1 = \begin{bmatrix} 3 \\ -3 \\ 0 \end{bmatrix}$ ,  $u^2 = \begin{bmatrix} 2 \\ 2 \\ -1 \end{bmatrix}$ ,  $u^3 = \begin{bmatrix} 1 \\ 1 \\ 4 \end{bmatrix}$ ,  $b = \begin{bmatrix} 5 \\ -3 \\ 1 \end{bmatrix}$ .

a. Form the matrix  $U = [u^1 \ u^2 \ u^3]$  and confirm that the columns of  $U$  are orthogonal by computing  $U^T U$ .

$$U = \begin{bmatrix} 3 & 2 & 1 \\ -3 & 2 & 1 \\ 0 & -1 & 4 \end{bmatrix}, U^T U = \begin{bmatrix} 3 & -3 & 0 \\ 2 & 2 & -1 \\ 1 & 1 & 4 \end{bmatrix} \begin{bmatrix} 3 & 2 & 1 \\ -3 & 2 & 1 \\ 0 & -1 & 4 \end{bmatrix} = \begin{bmatrix} 18 & 0 & 0 \\ 0 & 9 & 0 \\ 0 & 0 & 18 \end{bmatrix} \text{ which is}$$

diagonal so the columns of  $U$  are orthogonal.

b. Express  $b$  as a linear combination of  $u^1, u^2$  and  $u^3$ . (That is, solve  $Ux = b$ . Be clever about using  $U^T$  to do this.)

$$\text{If } Ux = b, \text{ then } U^T Ux = U^T b, \text{ so } \begin{bmatrix} 18 & 0 & 0 \\ 0 & 9 & 0 \\ 0 & 0 & 18 \end{bmatrix} x = \begin{bmatrix} 3 & -3 & 0 \\ 2 & 2 & -1 \\ 1 & 1 & 4 \end{bmatrix} \begin{bmatrix} 5 \\ -3 \\ 1 \end{bmatrix} = \begin{bmatrix} 24 \\ 3 \\ 6 \end{bmatrix}, \text{ so}$$

$$x = \begin{bmatrix} 24/18 \\ 3/9 \\ 6/18 \end{bmatrix} = \begin{bmatrix} 4/3 \\ 1/3 \\ 1/3 \end{bmatrix}.$$

2. Let  $u^1 = \begin{bmatrix} -2/3 \\ 1/3 \\ 2/3 \end{bmatrix}$ ,  $u^2 = \begin{bmatrix} 1/3 \\ 2/3 \\ 0 \end{bmatrix}$

a. Form the matrix  $U = [u^1 \ u^2]$  and confirm that the columns of  $U$  are orthogonal by computing  $U^T U$ .

$$U = \begin{bmatrix} -2/3 & 1/3 \\ 1/3 & 2/3 \\ 2/3 & 0 \end{bmatrix}, U^T U = \begin{bmatrix} -2/3 & 1/3 & 2/3 \\ 1/3 & 2/3 & 0 \end{bmatrix} \begin{bmatrix} -2/3 & 1/3 \\ 1/3 & 2/3 \\ 2/3 & 0 \end{bmatrix} = \begin{bmatrix} 9/9 & 0 \\ 0 & 5/9 \end{bmatrix}$$

which is diagonal so the columns of  $U$  are orthogonal.

b. Normalize the columns and confirm that  $U^T U = I$ .

After normalization,

$$U = \begin{bmatrix} -2/3 & 1/\sqrt{5} \\ 1/3 & 2/\sqrt{5} \\ 2/3 & 0 \end{bmatrix}, U^T U = \begin{bmatrix} -2/3 & 1/3 & 2/3 \\ 1/\sqrt{5} & 2/\sqrt{5} & 0 \end{bmatrix} \begin{bmatrix} -2/3 & 1/\sqrt{5} \\ 1/3 & 2/\sqrt{5} \\ 2/3 & 0 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}.$$

3. Answer true or false to the following. If false offer a counterexample.

a. Every orthogonal set in  $\mathbb{R}^n$  is linearly independent.

**False.** The set consisting of a zero vector alone is orthogonal but not linearly independent.

b. If a set  $S = \{u^1, u^2, \dots, u^k\}$  has the property that  $u^i \cdot u^j = 0$  whenever  $i \neq j$ , then  $S$  is an orthonormal set.

**False.** The set consisting of a zero vector alone has the property that  $u^i \cdot u^j = 0$  whenever  $i \neq j$ , but  $S$  is not an orthonormal set.

4. Show that if  $U$  is a square matrix whose columns are orthonormal then  $U^T = U^{-1}$ .

We have  $U^T U = I$ , so  $U^T = U^{-1}$ .

5. Show that if  $U$  is an  $m \times n$  orthogonal matrix then for all  $x \in \mathbb{R}^n$ ,  $\|Ux\| = \|x\|$ . (This is not hard: work out  $\|Ux\|^2$ . This can be stated as "An orthogonal transformation preserves length.")

$$\text{For all } x \in \mathbb{R}^n, \|Ux\|^2 = (Ux)^T (Ux) = x^T U^T U x = x^T x = \|x\|^2.$$

6. Consider this mathematical (and not necessarily computer) procedure:

$$[\alpha, v'] = \text{project}[u, v]$$

Inputs vectors  $u$  and  $v$ , computes and Returns  $\alpha = u \cdot v / u \cdot u$  and  $v' = v - \alpha u$ .

$$\text{Now, let } u^1 = \begin{bmatrix} 3 \\ -3 \\ 0 \end{bmatrix}, u^2 = \begin{bmatrix} -1 \\ 5 \\ -1 \end{bmatrix}, u^3 = \begin{bmatrix} 9 \\ -3 \\ 3 \end{bmatrix}.$$

a.  $[r_{1,2}, u_2'] = \text{project}[u_1, u_2]$ . (That is, subtract the projection of  $u_2$  onto the subspace spanned by  $u_1$ .)

$$\text{project} \left[ \begin{bmatrix} 3 \\ -3 \\ 0 \end{bmatrix}, \begin{bmatrix} -1 \\ 5 \\ -1 \end{bmatrix} \right] \text{ yields } r_{1,2} = -18/18 = -1, \text{ and } u_2' = \begin{bmatrix} -1 \\ 5 \\ -1 \end{bmatrix} - (-1) \begin{bmatrix} 3 \\ -3 \\ 0 \end{bmatrix} = \begin{bmatrix} 2 \\ 2 \\ -1 \end{bmatrix}.$$

b.  $[r_{1,3}, u_3'] = \text{project}[u_1, u_3]$ . (That is, subtract the projection of  $u_3$  onto the subspace spanned by  $u_1$ .)

$$\text{project} \left[ \begin{bmatrix} 3 \\ -3 \\ 0 \end{bmatrix}, \begin{bmatrix} 9 \\ -3 \\ 3 \end{bmatrix} \right] \text{ yields } r_{1,3} = 36/18 = 2, \text{ and } u_3' = \begin{bmatrix} 9 \\ -3 \\ 3 \end{bmatrix} - 2 \begin{bmatrix} 3 \\ -3 \\ 0 \end{bmatrix} = \begin{bmatrix} 3 \\ 3 \\ 3 \end{bmatrix}.$$

c.  $[r_{2,3}, u_3''] = \text{project}[u_2', u_3']$ . (That is, subtract the projection of  $u_3'$  onto the subspace spanned by  $u_2'$ .)

$$\text{project} \left[ \begin{bmatrix} 2 \\ 2 \\ -1 \end{bmatrix}, \begin{bmatrix} 3 \\ 3 \\ 3 \end{bmatrix} \right] \text{ yields } r_{2,3} = 9/9 = 1, \text{ and } u_3'' = \begin{bmatrix} 3 \\ 3 \\ 3 \end{bmatrix} - 1 \begin{bmatrix} 2 \\ 2 \\ -1 \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \\ 4 \end{bmatrix}.$$

d. Compute  $A = [u_1 \ u_2 \ u_3] \begin{bmatrix} 1 & r_{1,2} & r_{1,3} \\ 0 & 1 & r_{2,3} \\ 0 & 0 & 1 \end{bmatrix}$ . (Compare to  $U$  in Problem 1. You have

just used the Gram-Schmidt Algorithm to orthogonalize - but not **orthonormalize** - vectors. That is, the normalizations are not done.)

$$A = \begin{bmatrix} 3 & 2 & 1 \\ -3 & 2 & 1 \\ 0 & -1 & 4 \end{bmatrix} \begin{bmatrix} 1 & -1 & 2 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 3 & -1 & 9 \\ -3 & 5 & -3 \\ 0 & -1 & 3 \end{bmatrix} = [u_1 \ u_2 \ u_3] \text{ and}$$

$$[u_1 \ u_2 \ u_3] = \begin{bmatrix} 3 & 2 & 1 \\ -3 & 2 & 1 \\ 0 & -1 & 4 \end{bmatrix} = U \text{ in Problem 1.}$$

7. Prove that if  $y^T x = 0$ , for all  $x$ , then  $y = 0$ .  $y = 0$ . (Hint: Consider  $x = y$ , in particular.)

Since  $y^T x = 0$ , for all  $x$ , then in particular for  $x = y$ ,  $\|y\|^2 = y^T y = 0$ , and if  $\|y\| = 0$ ,  $y = 0$ .

8. Prove that if  $Q^T Q = I$ , then if  $x$  is perpendicular to  $y$ , then  $Qx$  is perpendicular to  $Qy$ .

If  $x^T y = 0$ , then  $(Qx)^T (Qy) = x^T Q^T Q y = x^T y = 0$ .