


# CS 343H: Honors Artificial Intelligence

## Bayes Nets: Independence



Prof. Peter Stone — The University of Texas at Austin

# Probability Recap

- Conditional probability

$$P(x|y) = \frac{P(x, y)}{P(y)}$$

- Product rule

$$P(x, y) = P(x|y)P(y)$$

- Chain rule

$$\begin{aligned} P(X_1, X_2, \dots, X_n) &= P(X_1)P(X_2|X_1)P(X_3|X_1, X_2) \dots \\ &= \prod_{i=1}^n P(X_i|X_1, \dots, X_{i-1}) \end{aligned}$$

- X, Y independent if and only if:

$$\forall x, y : P(x, y) = P(x)P(y)$$

- X and Y are conditionally independent given Z if and only if:

$$\forall x, y, z : P(x, y|z) = P(x|z)P(y|z)$$

$$X \perp\!\!\!\perp Y | Z$$

# Bayes Nets

- A Bayes' net is an efficient encoding of a probabilistic model of a domain



- Questions we can ask:
  - Inference: given a fixed BN, what is  $P(X | e)$ ?
  - Representation: given a BN graph, what kinds of distributions can it encode?
  - Modeling: what BN is most appropriate for a given domain?

# Bayes Net Semantics


- A directed, acyclic graph, one node per random variable
- A conditional probability table (CPT) for each node
  - A collection of distributions over  $X$ , one for each combination of parents' values

$$P(X|a_1 \dots a_n)$$

- Bayes' nets implicitly encode joint distributions


- As a product of local conditional distributions
- To see what probability a BN gives to a full assignment, multiply all the relevant conditionals together:

$$P(x_1, x_2, \dots, x_n) = \prod_{i=1}^n P(x_i | \text{parents}(X_i))$$



# Example: Alarm Network

B	P(B)
+b	0.001
-b	0.999



E	P(E)
+e	0.002
-e	0.998

A	M	P(M A)
+a	+m	0.7
+a	-m	0.3
-a	+m	0.01
-a	-m	0.99




A	J	P(J A)
+a	+j	0.9
+a	-j	0.1
-a	+j	0.05
-a	-j	0.95

B	E	A	P(A B,E)
+b	+e	+a	0.95
+b	+e	-a	0.05
+b	-e	+a	0.94
+b	-e	-a	0.06
-b	+e	+a	0.29
-b	+e	-a	0.71
-b	-e	+a	0.001
-b	-e	-a	0.999

$$P(+b, -e, +a, -j, +m) =$$

# Example: Alarm Network

B	P(B)
+b	0.001
-b	0.999



E	P(E)
+e	0.002
-e	0.998

A	M	P(M A)
+a	+m	0.7
+a	-m	0.3
-a	+m	0.01
-a	-m	0.99



A	J	P(J A)
+a	+j	0.9
+a	-j	0.1
-a	+j	0.05
-a	-j	0.95

B	E	A	P(A B,E)
+b	+e	+a	0.95
+b	+e	-a	0.05
+b	-e	+a	0.94
+b	-e	-a	0.06
-b	+e	+a	0.29
-b	+e	-a	0.71
-b	-e	+a	0.001
-b	-e	-a	0.999

$$\begin{aligned}
 &P(+b, -e, +a, -j, +m) = \\
 &P(+b)P(-e)P(+a|+b, -e)P(-j|+a)P(+m|+a) = \\
 &0.001 \times 0.998 \times 0.94 \times 0.1 \times 0.7
 \end{aligned}$$

# Size of a Bayes Net

- How big is a joint distribution over  $N$  Boolean variables?

$$2^N$$



- How big is an  $N$ -node net if nodes have up to  $k$  parents?

$$O(N * 2^{k+1})$$

- Both give you the power to calculate

$$P(X_1, X_2, \dots, X_n)$$

- BNs: Huge space savings!
- Also easier to elicit local CPTs
- Also faster to answer queries (coming)



# Bayes Nets

---

- ✓ Representation
  - Conditional Independences
  - Probabilistic Inference
  - Learning Bayes' Nets from Data



# Conditional Independence

- X and Y are **independent** if

$$\forall x, y \quad P(x, y) = P(x)P(y) \quad \text{---} \rightarrow \quad X \perp\!\!\!\perp Y$$

- X and Y are **conditionally independent** given Z

$$\forall x, y, z \quad P(x, y|z) = P(x|z)P(y|z) \quad \text{---} \rightarrow \quad X \perp\!\!\!\perp Y|Z$$

- (Conditional) independence is a property of a distribution

- Example:  $Alarm \perp\!\!\!\perp Fire|Smoke$



# Bayes Nets: Assumptions


- Assumptions we are required to make to define the Bayes net when given the graph:

$$P(x_i | x_1 \cdots x_{i-1}) = P(x_i | \text{parents}(X_i))$$

- Beyond above “chain rule → Bayes net” conditional independence assumptions:
  - Often additional conditional independences
  - They can be inferred from the graph structure
- Important for modeling: understand assumptions made when choosing a Bayes net graph



# Example



- Conditional independence assumptions directly from simplifications in chain rule:

Standard chain rule:  $p(x, y, z, w) = p(x)p(y|x)p(z|x, y)p(w|x, y, z)$


Bayes net:  $p(x, y, z, w) = p(x)p(y|x)p(z|y)p(w|z)$

Since:  $z \perp\!\!\!\perp x \mid y$  and  $w \perp\!\!\!\perp x, y \mid z$  (cond. indep. given parents)

- Additional implied conditional independence assumptions?  $w \perp\!\!\!\perp x \mid y$

$$\begin{aligned} p(w|x, y) &= \frac{p(w, x, y)}{p(x, y)} = \frac{\sum_z p(x)p(y|x)p(z|y)p(w|z)}{p(x)p(y|x)} = \sum_z p(z|y)p(w|z) = \sum_z p(z|y)p(w|z, y) \\ &= \sum_z p(z, w|y) = p(w|y) \end{aligned}$$

# D-separation: Outline



# D-separation: Outline

---

- Study independence properties for triples
- Analyze complex cases in terms of member triples
- D-separation: a condition / algorithm for answering such queries

# Causal Chains

- This configuration is a “causal chain”



X: Low pressure

Y: Rain

Z: Traffic

$$P(x, y, z) = P(x)P(y|x)P(z|y)$$

- Guaranteed X independent of Z ? *No!*

- One example set of CPTs for which X is not independent of Z is sufficient to show this independence is not guaranteed.

- Example:

- Low pressure causes rain causes traffic, high pressure causes no rain causes no traffic

- In numbers:

$$P(+y | +x) = 1, P(-y | -x) = 1, \\ P(+z | +y) = 1, P(-z | -y) = 1$$

# Causal Chains

- This configuration is a “causal chain”

- Guaranteed X independent of Z given Y?



X: Low pressure

Y: Rain

Z: Traffic

$$P(x, y, z) = P(x)P(y|x)P(z|y)$$


$$\begin{aligned} P(z|x, y) &= \frac{P(x, y, z)}{P(x, y)} \\ &= \frac{P(x)P(y|x)P(z|y)}{P(x)P(y|x)} \\ &= P(z|y) \end{aligned}$$

*Yes!*

- Evidence along the chain “blocks” the influence

# Common Cause

- This configuration is a “common cause”



$$P(x, y, z) = P(y)P(x|y)P(z|y)$$

- Guaranteed X independent of Z ? *No!*

- One example set of CPTs for which X is not independent of Z is sufficient to show this independence is not guaranteed.

- Example:

- Project due causes both forums busy and lab full

- In numbers:

$$P(+x \mid +y) = 1, P(-x \mid -y) = 1,$$


$$P(+z \mid +y) = 1, P(-z \mid -y) = 1$$



# Common Cause

- This configuration is a “common cause”

- Guaranteed X and Z independent given Y?



$$P(x, y, z) = P(y)P(x|y)P(z|y)$$


$$\begin{aligned} P(z|x, y) &= \frac{P(x, y, z)}{P(x, y)} \\ &= \frac{P(y)P(x|y)P(z|y)}{P(y)P(x|y)} \\ &= P(z|y) \end{aligned}$$

*Yes!*

- Observing the cause blocks influence between effects.

# Common Effect

- Last configuration: two causes of one effect (v-structures)



- Are X and Y independent?
  - **Yes**: the ballgame and the rain cause traffic, but they are not correlated
  - Still need to prove they must be (try it!)
- Are X and Y independent given Z?
  - **No**: seeing traffic puts the rain and the ballgame in competition as explanation.
- **This is backwards from the other cases**
  - Observing an effect **activates** influence between possible causes.


# The General Case

---




# The General Case

- General question: in a given BN, are two variables independent (given evidence)?
- Solution: analyze the graph
- Any complex example can be broken into repetitions of the three canonical cases



# Reachability




- Recipe: shade evidence nodes, look for paths in the resulting graph
- Attempt 1: if two nodes are connected by an undirected path not blocked by a shaded node, they are conditionally dependent
- Almost works, but not quite
  - Where does it break?
  - Answer: the v-structure at T doesn't count as a link in a path unless "active"






# Active / Inactive Paths

- Question: Are X and Y conditionally independent given evidence variables {Z}?
  - Yes, if X and Y “d-separated” by Z
  - Consider all (undirected) paths from X to Y
  - No active paths = independence!
- A path is active if each triple is active:
  - Causal chain  $A \rightarrow B \rightarrow C$  where B is unobserved (either direction)
  - Common cause  $A \leftarrow B \rightarrow C$  where B is unobserved
  - Common effect (aka v-structure)  
 $A \rightarrow B \leftarrow C$  where B or one of its descendants is observed
- All it takes to block a path is a single inactive segment

Active Triples



Inactive Triples




# D-Separation

- Query:  $X_i \perp\!\!\!\perp X_j \mid \{X_{k_1}, \dots, X_{k_n}\} ?$
- Check all (undirected!) paths between  $X_i$  and  $X_j$ 
  - If one or more active, then independence not guaranteed

$$X_i \not\perp\!\!\!\perp X_j \mid \{X_{k_1}, \dots, X_{k_n}\}$$

- Otherwise (i.e. if all paths are inactive), then independence is guaranteed

$$X_i \perp\!\!\!\perp X_j \mid \{X_{k_1}, \dots, X_{k_n}\}$$




# Example

---

$R \perp\!\!\!\perp B$       *Yes*

$R \perp\!\!\!\perp B | T$       *No*

$R \perp\!\!\!\perp B | T'$       *No*





# Example


$L \perp\!\!\!\perp T' \mid T$     *Yes*

$L \perp\!\!\!\perp B$     *Yes*

$L \perp\!\!\!\perp B \mid T$     *No*

$L \perp\!\!\!\perp B \mid T'$     *No*

$L \perp\!\!\!\perp B \mid T, R$     *Yes*



# Example

- Variables:


- R: Raining
- T: Traffic
- D: Roof drips
- S: I'm sad

- Questions:

$T \perp\!\!\!\perp D$  *No*

$T \perp\!\!\!\perp D | R$  *Yes*

$T \perp\!\!\!\perp D | R, S$  *No*




# Structure Implications

- Given a Bayes net structure, can run d-separation algorithm to build a complete list of conditional independences that are necessarily true of the form


$$X_i \perp\!\!\!\perp X_j \mid \{X_{k_1}, \dots, X_{k_n}\}$$

- This list determines the set of probability distributions that can be represented




# Computing All Independences


COMPUTE ALL THE  
INDEPENDENCES!




$$X \perp\!\!\!\perp Z \mid Y$$



$$X \perp\!\!\!\perp Z \mid Y$$



$$X \perp\!\!\!\perp Z$$







**None!**

# Topology Limits Distributions

- Given some graph topology  $G$ , only certain joint distributions can be encoded
- The graph structure guarantees certain (conditional) independences
- (There might be more independence)
- Adding arcs increases the set of distributions, but has several costs
- Full conditioning can encode any distribution

$$\{X \perp\!\!\!\perp Y, X \perp\!\!\!\perp Z, Y \perp\!\!\!\perp Z, \\ X \perp\!\!\!\perp Z \mid Y, X \perp\!\!\!\perp Y \mid Z, Y \perp\!\!\!\perp Z \mid X\}$$



# Bayes Nets Representation Summary

---

- Bayes nets compactly encode joint distributions
- Guaranteed independencies of distributions can be deduced from BN graph structure
- D-separation gives precise conditional independence guarantees from graph alone
- A Bayes net's joint distribution may have further (conditional) independence that is not detectable until you inspect its specific distribution

# Bayes Nets

---

✓ Representation

✓ Conditional Independences

- Probabilistic Inference

- Enumeration (exact, exponential complexity)
- Variable elimination (exact, worst-case exponential complexity, often better)
- Probabilistic inference is NP-complete
- Sampling (approximate)

- Learning Bayes' Nets from Data